



# A Study of Vegetation is an Important Indication data for the Distribution or Communities and their Relation with Corporeal Environment of Certain Area

**Ujevic Jha**

*Assistant Professor, Department of Economics, Indira Gandhi University, Meerpur, Haryana, India*

**Chakravarthy Darko**

*Assistant Professor, Department of Economics, Indira Gandhi University, Meerpur, Haryana, India*

**Dharam Hooda\***

*Student, Department of Economics, Indira Gandhi University, Meerpur, Haryana, India*

\*Corresponding author

## Abstract

Pteridophytes are the primitive vascular plants, they are found scattered all over the globe and quite many of them occur in India. However, they are not found throughout the country. Pteridophytes make an important contribution to the earth's plant diversity. Being the second largest group of vascular plants, they form a significant and dominant component of many plant communities. All human activities disturb the vegetation by eliminating many species. So an up to date study is very important for knowing about flora, especially in the case of pteridophytes. The present effort was undertaken to enumerate the pteridophytic species present in Kilavarai river situated in Palni hills of southern Western Ghats, Tamil Nadu, India, the study was carried out from December 2014 to January 2016. The result of the present study revealed that 36 pteridophytic species belonging to 25 genera distributed among 19 families were noted from the study site. Among the families recorded, Adiantaceae (16.67%) was found to be dominant and largest family comprising 6 species. Polypodiaceae and Pteridaceae (each of 11.11%) shared the second largest family status; they were represented by 4 species each. The third largest family was Lycopodiaceae (8.33%); it was represented by 3 species. The families Cheilantheaceae, and Selaginellaceae were recorded with each of 2 species (5.56%) and 11 families were represented by single species (2.78%). Moreover, terrestrial species (27 species) were found to be more in number than epiphytic (5 species), 3 species were found as both epiphytic and lithophytic species and 1 species was recorded as purely lithophytic.

**Keywords:** Pteridophytes, Kilavarai, Palni hills, Western Ghats, Nephrolepidaceae, Freshwater, Pteridophytes

## 1. Introduction

Pteridophytes may comprise a significant component of the forest ecosystem. Ferns have measurable indications that may reflect the effects of change in environmental factors. The ferns are not only taxonomic oddities but those are plants with dynamic relationship to their environment (Verma and Khullar, 2010). Ecological study of flora is an important indication as it forms baseline data for the distribution of plant species or communities and their relation with physical environment of particular area (Birand Vasudeva, 1972). Western Ghats are very rich and varied in flora because of its diversified topography and varied climatic conditions. The region is considered to be one of 34 major hot spots of biological diversity in the world (Myers, 2003). Though the pteridophytes occur in abundance in the tropical, sub-tropical and moist deciduous forests of India, large scale destruction of forests has drastically affected the diversity of pteridophyte species.

The life-cycle of the ferns and fern-allies is dependent upon the existence of forests, but due to habitat destruction many species have been reduced and the rare ferns are being extinct or are on the verge of extinction (Dixit, 2000). A survey of literature shows that areas rich in the diversity of plants including the angiosperms to the lower group such as pteridophytes, bryophytes, lichens, fungi and algae. But the works carried out with the lower group of plants are very limited. However, pteridophytic plants have not received sufficient attention by plant explorers of forest of upper Palani hills due to difficulties in the species identification. Therefore, present work

deals with the distribution of pteridophytes in Kilavarai freshwater river situated in the upper Palni hills of Southern Western Ghats, Dindigul district, Tamilnadu, India.

## 2. Materials and Methods

### 2.1 Study Area

The present study was carried out in Kilavarai freshwater river, located in the upper Palni hills of Kodaikanal situated in Southern Palni hills (10° 23' 89.62"N and 77° 31' 50.69"E) of Western Ghats, Dindigul district, TamilNadu, India. This perennial river develops into Vanderavu huts flows west upto Amaravathy reservoir. The flowing area of the river, with varying degree of slopes leading into short and height waterfalls, like Polur waterfalls. Kilavarai river is located along Kilavarai –Polur –Kumbur villages path. The experimental river has been utilized for domestic and agricultural purposes of the inhabitants of near and around the villages.

### 2.2 Survey And identification

An extensive and intensive floristic survey of the pteridophyte was carried out during December 2014 to January 2016, by employing collection, identification and verification. Specimens of pteridophytic plants found in the study area were collected and processed in the laboratory. The specimens were poisoned with mercuric chloride. Herbaria for the plant specimens which have been prepared and deposited in the Post Graduate and Research Department of Botany, Saraswathi Narayanan College, Madurai. The pteridophytic plants were identified using standard keys and flora: The Ferns of Southern India Madras: A handbook to the ferns of British India Ceylon and Malay Peninsula and The Ferns of British India by R.H. Beddome (1873, 1892 & 1865-1870), Fern Flora of the Palni Hills (South India) by V.S. Manickam (1986), Pteridophyte Flora of the Western Ghats –South India by Manickam and Irudayaraj, (1992), Pteridophyte Flora of Nilgiris, South India (Manickam and Irudayaraj, (2003) and through comparison with the online floras. The documented species were arranged based on Pichi-Sermolli's (1977) system for the present systematic treatment.

## 3. Result and Discussion

The present survey reveals that a total of 36 pteridophytic species belonging to 25 genera distributed among 19 families were documented (Table 1) from the study area. Among the pteridophytic species recorded, Adiantum was represented by 6 species and which was found to be the dominant genus, and it was followed by the genus, Pteris with 3 species. A total of 5 genera viz., Cheilanthes, Christella, Nephrolepis, Pyrrosia and Selaginella were represented with each of 2 species and the rest of the 17 genus were represented as monospecific genus. Among the families recorded, Adiantaceae (16.67%) was found to be dominant and largest family comprising 6 species.

**Table 1:** Pteridophytic diversity of Kilavarai freshwater river. (Terrestrial (T), Epiphyte (E), Lithophyte (L))

1.	<i>Huperzia phlegmaria</i> (L.) Holub	Lycopodiaceae	E
2.	<i>Lycopodiella cernua</i> (L.) Pic.	Lycopodiaceae	T
3.	<i>Lycopodium clavatum</i> auct. Clarke	Lycopodiaceae	T
4.	<i>Selaginella involvens</i> (Sw.) Spring	Selaginellaceae	E
5.	<i>Selaginella wightii</i> Hieron	Selaginellaceae	E
6.	<i>Equisetum ramosissimum</i> Desr.	Equisetaceae	T
7.	<i>Angiopteris evecta</i> (G. Forst.) Hoffm.	Angiopteridaceae	T
8.	<i>Osmunda hugeliana</i> Presl.	Osmundaceae	T
9.	<i>Dicranopteris linearis</i> (Burm.f.) Underwood,	Glecheniaceae	T
10.	<i>Drynaria quercifolia</i> (L.) J. Sm	Polypodiaceae	E
11.	<i>Lepisorus nudus</i> (Hook.) Ching	Polypodiaceae	E / L
12.	<i>Pyrrosia lanceolata</i> Farewell	Polypodiaceae	E / L
13.	<i>Pyrrosia porosa</i> (C.Presl) Hovenkamp	Polypodiaceae	E / L
14.	<i>Doryopteris concolor</i> (Langsd. Et Fisch.) Kuhn	Sinopteridaceae	L
15.	<i>Cheilanthes bulbosa</i> Kuntze	Cheilanthaceae	T
16.	<i>Cheilanthes mysorensis</i> Wall.	Cheilanthaceae	T
17.	<i>Pteridium aquilinum</i> (L.) Kuhn	Pteridaceae	T
18.	<i>Pteris biaurita</i> L.	Pteridaceae	T
19.	<i>Pteris confusa</i> T.G. Walker	Pteridaceae	T
20.	<i>Pteris gongalensis</i> T.G. Walker	Pteridaceae	T
21.	<i>Actinopteris radiata</i> (Sw.) Link.	Actiniopteridaceae	E
22.	<i>Adiantum capillus-veneris</i> L.	Adiantaceae	T
23.	<i>Adiantum hispidulum</i> Sw.	Adiantaceae	T
24.	<i>Adiantum incisum</i> Forssk.	Adiantaceae	T
25.	<i>Adiantum lunulatum</i> Burm.f.	Adiantaceae	T
26.	<i>Adiantum raddianum</i> C. Presl.	Adiantaceae	T

Polypodiaceae and Pteridaceae (each of 11.11%) shared the second largest family status; they were represented by 4 species each. The third largest family was Lycopodiaceae (8.33%); it was represented by 3 species. The families Cheilantheaceae, Nephrolepidaceae and Selaginellaceae were recorded with each of 2 species (5.56%) and 11 families (Actiniopteridaceae, Angiopteridaceae, Aspleninaceae, Cyatheaceae, Dryopteridaceae, Equisetaceae, Gleicheniaceae, Hemionitidaceae, Lindsaceae, Osmundaceae and Sinopteridaceae) were represented by single species (2.78%) (Table 1; Figure 1). The present study also indicated that during rainy season, all kinds of plants including ferns exhibited a very luxurious growth. Similar observation is also supported by many reports on pteridophytes of the Western Ghats (Sukumaran et al., 2006). Species diversity of pteridophytes slowly get decreased and today they are restricted to lesser habitat.

Regarding the habitats of the pteridophytic species recorded by the present investigation, terrestrial species (27 species) were found to be more in number than epiphytic (5 species), 3 species were found as both epiphytic and lithophytic species and 1 species was recorded as purely lithophytic (Figure 2). These results were corroborated with the findings of a study carried out in a miniature sacred groves of Kanyakumari district to document the pteridophytic flora from the same by Sukumaran et al. (2009) and they have found that the terrestrial pteridophytic species more in number than epiphytic and lithophytic species.




Figure 2: Habitat of Pteridophytic species.

#### 4. Conclusion

The conservation focus should not be restricted upto the commercially important species and specific strategies for conservation of pteridophytes should be taken into account with much consideration. Botanical explorations should increase in the under-explored botanically rich areas for documenting the diversity and ecological characteristics of pteridophytes and taxonomic reinvestigations should take place in order to avoid the confusions with new species and existing species.

#### Reference

- Alagesabooopathi, C and G. Subramanian. (2017). Ethnomedicinal studies on some Pteridophytes of Kanjamalai Hills, Salem district of Tamilnadu, India. *World Journal of Pharmaceutical Research*, 6(8): 1590-1596.
- Beddome, R.H. (1863-1865). *The Ferns of Southern India*. Gantz Brothers, Madras.
- Beddome, R.H. (1865-1870). *The Ferns of British India*. Vol. 1 and 2. Gantz Bros, Madras (Reprinted.1973). Oxford and IBH Publishing Co., New Delhi.
- Beddome, R.H. (1883). *A Hand book to the Ferns of British India, Ceylon, and Malay Peninsula*. Thacker Spink and Co., Calcutta ((Reprint ed. 1976). Today and Tomorrow's Printers, New Delhi.
- Bir, S.S and Vasudeva, S.M. (1972). Ecological and Phytogeographical observation on the Pteridophytic flora of Pachmarhi Hills (Central India). *J. Indian Bot. Soc*, 15: 297-304.
- Dixit, R.D. (2000). Conspectus of Pteridophytic diversity in India. *Indian Fern Journal*, 17: 77 -91.
- Manickam V. S and V. Irudayaraj. (1992). *Pteridophyte Flora of the Western Ghats-South India*, BI Publications Pvt Ltd. New Delhi.
- Manickam, V.S. (1986). *Fern Flora of the Palni Hills (South India)*. Today and Tomorrow's Printers and Publishers, New Delhi.
- Manickam, V.S. and V. Irudayaraj. (2003). *Pteridophyte Flora of Nilgiris, South India*, Bishen Singh Mahendra Pal Singh, Dehra Dun
- Myers, N. (2003). Biodiversity revisited. *Biosciences*, 53: 916-917.

- 11.Sukumaran, S, Jeeva, Sand A.D.S. Raj. (2009). Diversity of pteridophytes in miniature sacred forests of Kanyakumari district, Southern Western Ghats.Indian J Forestry,32(2): 285-290.
- 12.Verma, S.C and S. P. Khullar. (2010). Book Review on Fern Ecology.Indian Fern Journal,27(1-2): 383-387.

